



It's important that you return for your next dose as directed by your doctor

SIGN UP

Prolia.com/bonematters
Or Call 1-866-426-6355

Please see Important Safety Information inside

Amgen Inc., One Amgen Center Drive, Thousand Oaks, CA 91320-1799
© 2019 Amgen Inc. All rights reserved. USA-162-81634 11/19



For women with postmenopausal osteoporosis with high risk for fracture and for men with osteoporosis with high risk for fracture



Bone Matters®: An Empowering Patient Support Program

Sign up to receive:



Welcome kit to get you started



Lifestyle information, including exercise tips, calcium-rich recipes, and more



Courtesy reminders* to help keep you on track for your next Prolia® injection



Call center access for questions or to update your treatment appointments

*This program is intended for the limited purpose of delivering courtesy reminder notifications to remind you to schedule your next injection of Prolia®. However, it is up to you and/or your caregiver to ensure you obtain your next injection in a timely manner and in accordance with your doctor's instruction. Please go to www.prolia.com for the full Program details and information.

What is Prolia®? Prolia® is a prescription medicine used to treat osteoporosis in women after menopause who are at high risk for fracture or cannot use another osteoporosis medicine or other osteoporosis medicines did not work well. Prolia® is a prescription medicine used to increase bone mass in men with osteoporosis who are at high risk for fracture.

Please see Important Safety Information inside

Indications and Important Safety Information

Indications:

Prolia® is a prescription medicine used to treat osteoporosis in women after menopause who are at high risk for fracture or cannot use another osteoporosis medicine or other osteoporosis medicines did not work well.

Prolia® is a prescription medicine used to increase bone mass in men with osteoporosis who are at high risk for fracture.

Important Safety Information

Do not take Prolia® if you: have low blood calcium; or are pregnant or plan to become pregnant, as Prolia® may harm your unborn baby; or are allergic to denosumab or any ingredients in Prolia®.

What is the most important information I should know about Prolia®?

If you receive Prolia®, you should not receive XGEVA®. Prolia® contains the same medicine as XGEVA® (denosumab).

Prolia® can cause serious side effects:

Serious allergic reactions have happened in people who take Prolia®. Call your doctor or go to your nearest emergency room right away if you have any symptoms of a serious allergic reaction, including low blood pressure (hypotension); trouble breathing; throat tightness; swelling of your face, lips, or tongue; rash; itching; or hives.

Low blood calcium (hypocalcemia). Prolia® may lower the calcium levels in your blood. If you have low blood calcium, it may get worse during treatment. Your low blood calcium must be treated before you receive Prolia®.

Take calcium and vitamin D as your doctor tells you to help prevent low blood calcium.

Severe jaw bone problems (osteonecrosis) may occur. Your doctor should examine your mouth before you start Prolia® and may tell you to see your dentist. It is important for you to practice good mouth care during treatment with Prolia®.

Unusual thigh bone fractures. Some people have developed unusual fractures in their thigh bone. Symptoms of a fracture include new or unusual pain in your hip, groin, or thigh.

Increased risk of broken bones, including broken bones in the spine, after stopping, skipping or delaying Prolia®. Talk with your doctor before starting Prolia® treatment. After your treatment with Prolia® is stopped, or if you skip or delay taking a dose, your risk for breaking bones, including bones in your spine, is increased. Your risk for having more than 1 broken bone in your spine is increased if you have already had a broken bone in your spine. Do not stop, skip or delay taking Prolia® without first talking with your doctor. If your Prolia® treatment is stopped, talk to your doctor about other medicine that you can take.

Serious infections in your skin, lower stomach area (abdomen), bladder, or ear may happen. Inflammation of the inner lining of the heart (endocarditis) due to an infection may also happen more often in people who take Prolia®. You may need to go to the hospital for treatment.

Prolia® is a medicine that may affect the ability of your body to fight infections. People who have weakened immune systems or take medicines that affect the immune system may have an increased risk for developing serious infections.

Skin problems such as inflammation of your skin (dermatitis), rash, and eczema have been reported.

Bone, joint, or muscle pain. Some people who take Prolia® develop severe bone, joint, or muscle pain.

Before taking Prolia®, tell your doctor about all of your medical conditions, including if you:

- Take the medicine XGEVA® (denosumab)
- Have low blood calcium
- Cannot take daily calcium and vitamin D
- Had parathyroid or thyroid surgery (glands located in your neck)
- Have been told you have trouble absorbing minerals in your stomach or intestines (malabsorption syndrome)
- Have kidney problems or are on kidney dialysis
- Are taking medicine that can lower your blood calcium levels
- Plan to have dental surgery or teeth removed
- Are pregnant or plan to become pregnant

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with Prolia®.
- You should use an effective method of birth control (contraception) during treatment with Prolia® and for at least 5 months after your last dose of Prolia®.
- Tell your doctor right away if you become pregnant while taking Prolia®.
- Are breast-feeding or plan to breast-feed

What are the possible side effects of Prolia®?

It is not known if the use of Prolia® over a long period of time may cause slow healing of broken bones. The most common side effects of Prolia® in women being treated for osteoporosis after menopause are back pain, pain in your arms and legs, high cholesterol, muscle pain, and bladder infection. The most common side effects of Prolia® in men with osteoporosis are back pain, joint pain, and common cold (runny nose or sore throat).

These are not all the possible side effects of Prolia®. Call your doctor for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see accompanying Prolia® full Prescribing Information, including Medication Guide.